

# Hyperglycemic Crises in Patients With Diabetes Mellitus

AMERICAN DIABETES ASSOCIATION

**K**etoacidosis and hyperosmolar hyperglycemia are the two most serious acute metabolic complications of diabetes, even if managed properly. These disorders can occur in both type 1 and type 2 diabetes. The mortality rate in patients with diabetic ketoacidosis (DKA) is <5% in experienced centers, whereas the mortality rate of patients with hyperosmolar hyperglycemic state (HHS) still remains high at ~15%. The prognosis of both conditions is substantially worsened at the extremes of age and in the presence of coma and hypotension (1–10).

This position statement will outline precipitating factors and recommendations for the diagnosis, treatment, and prevention of DKA and HHS. It is based on the accompanying technical review (11), which should be consulted for further information.

**PATHOGENESIS** — Although the pathogenesis of DKA is better understood than that of HHS, the basic underlying mechanism for both disorders is a reduction in the net effective action of circulating insulin coupled with a concomitant elevation of counterregulatory hormones, such as glucagon, catecholamines, cortisol, and growth hormone. These hormonal alterations in DKA and HHS lead to increased hepatic and renal glucose production and impaired glucose utilization in peripheral tissues, which result in hyperglycemia and parallel changes in osmolality of the extracellular space (12,13). The combination of insulin deficiency and increased counterregulatory hormones in DKA also leads to release of free fatty acids into the circulation from adipose tissue (lipolysis) and to unre-

strained hepatic fatty acid oxidation to ketone bodies ( $\beta$ -hydroxybutyrate [ $\beta$ -OHB] and acetoacetate), with resulting ketonemia and metabolic acidosis. HHS on the other hand may be due to plasma insulin concentration inadequate to facilitate glucose utilization by insulin-sensitive tissues but adequate (as determined by residual C-peptide) to prevent lipolysis and subsequent ketogenesis, although the evidence for this is weak (14). Both DKA and HHS are associated with glycosuria, leading to osmotic diuresis with loss of water, sodium, potassium, and other electrolytes (3,15–20). The laboratory and clinical characteristics of DKA and HHS are summarized in Tables 1 and 2. As can be seen, DKA and HHS differ in magnitude of dehydration and degree of ketosis (and acidosis).

**PRECIPITATING FACTORS** — The most common precipitating factor in the development of DKA or HHS is infection. Other precipitating factors include cerebrovascular accident, alcohol abuse, pancreatitis, myocardial infarction, trauma, and drugs. In addition, newly onset type 1 diabetes or discontinuation of or inadequate insulin in established type 1 diabetes commonly leads to the development of DKA. Elderly individuals with newly onset diabetes (particularly residents of chronic care facilities) or individuals with known diabetes who become hyperglycemic and are unaware of it or are unable to take fluids when necessary are at risk for HHS (6).

Drugs that affect carbohydrate metabolism, such as corticosteroids, thiazides, and sympathomimetic agents (e.g., dobutamine and terbutaline), may precipitate the devel-

opment of HHS or DKA. In young patients with type 1 diabetes, psychological problems complicated by eating disorders may be a contributing factor in 20% of recurrent ketoacidosis. Factors that may lead to insulin omission in younger patients include fear of weight gain with improved metabolic control, fear of hypoglycemia, rebellion from authority, and stress of chronic disease (13).

## DIAGNOSIS

### History and physical examination

The process of HHS usually evolves over several days to weeks, whereas the evolution of the acute DKA episode in type 1 diabetes or even in type 2 diabetes tends to be much shorter. Although the symptoms of poorly controlled diabetes may be present for several days, the metabolic alterations typical of ketoacidosis usually evolve within a short time frame (typically <24 h). Occasionally, the entire symptomatic presentation may evolve or develop more acutely, and the patient may present in DKA with no prior clues or symptoms. For both DKA and HHS, the classical clinical picture includes a history of polyuria, polydipsia, polyphagia, weight loss, vomiting, abdominal pain (only in DKA), dehydration, weakness, clouding of sensoria, and finally coma. Physical findings may include poor skin turgor, Kussmaul respirations (in DKA), tachycardia, hypotension, alteration in mental status, shock, and ultimately coma (more frequent in HHS). Up to 25% of DKA patients have emesis, which may be coffee-ground in appearance and guaiac positive. Endoscopy has related this finding to the presence of hemorrhagic gastritis. Mental status can vary from full alertness to profound lethargy or coma, with the latter more frequent in HHS. Although infection is a common precipitating factor for both DKA and HHS, patients can be normothermic or even hypothermic primarily because of peripheral vasodilation. Hypothermia, if present, is a poor prognostic sign (21). Caution needs to be taken with patients who complain of abdominal pain on presentation, because the symptoms could be either a result or a cause (particularly in younger patients) of DKA. Further evaluation is necessary if this complaint does not resolve

.....

The recommendations in this paper are based on the evidence reviewed in the following publication: Management of hyperglycemic crises in patients with diabetes (Technical Review). *Diabetes Care* 24:131–153, 2001.

The initial draft of this position statement was prepared by Abbas E. Kitabchi, PhD, MD, Guillermo E. Umpierrez, MD, Mary Beth Murphy, RN, MS, CDE, MBA, Eugene J. Barrett, MD, PhD, Robert A. Kreisberg, MD, John I. Malone, MD, and Barry M. Wall, MD. The paper was peer-reviewed, modified, and approved by the Professional Practice Committee and the Executive Committee, October 2000.

**Abbreviations:**  $\beta$ -OHB,  $\beta$ -hydroxybutyric acid; AKA, alcoholic ketoacidosis; DKA, diabetic ketoacidosis; HHS, hyperosmolar hyperglycemic state.

**Table 1—Diagnostic criteria for DKA and HHS**

	DKA			HHS
	Mild	Moderate	Severe	
Plasma glucose (mg/dl)	>250	>250	>250	>600
Arterial pH	7.25–7.30	7.00–<7.24	<7.00	>7.30
Serum bicarbonate (mEq/l)	15–18	10–<15	<10	>15
Urine ketones*	Positive	Positive	Positive	Small
Serum ketones*	Positive	Positive	Positive	Small
Effective serum osmolality (mOsm/kg)†	Variable	Variable	Variable	>320
Anion gap‡	>10	>12	>12	<12
Alteration in sensoria or mental obtundation	Alert	Alert/drowsy	Stupor/coma	Stupor/Coma

\*Nitroprusside reaction method; †calculation:  $2[\text{measured Na (mEq/l)}] + \text{glucose (mg/dl)}/18$ ; ‡calculation:  $(\text{Na}^+) - (\text{Cl}^- + \text{HCO}_3^-)$  (mEq/l). See text for details.

with resolution of dehydration and metabolic acidosis.

### Laboratory findings

The initial laboratory evaluation of patients with suspected DKA or HHS should include determination of plasma glucose, blood urea nitrogen/creatinine, serum ketones, electrolytes (with calculated anion gap), osmolality, urinalysis, urine ketones by dipstick, as well as initial arterial blood gases, complete blood count with differential, and electrocardiogram. Bacterial cultures of urine, blood, and throat, etc., should be obtained and appropriate antibiotics given if infection is suspected.  $\text{HbA}_{1c}$  may be useful in determining whether this acute episode is the culmination of an evolutionary process in previously undiagnosed or poorly controlled diabetes or a truly acute episode in an otherwise well-controlled patient. A chest X ray should also be obtained if indicated. Tables 1 and 2 depict typical laboratory findings in patients with DKA or HHS.

The majority of patients with hyperglycemic emergencies present with leukocytosis proportional to blood ketone body concentration. Serum sodium concentration is usually decreased because of the osmotic flux of water from the intracellular to the extracellular space in the presence of hyperglycemia, and less commonly, serum sodium concentration may be falsely lowered by severe hypertriglyceridemia. Serum potassium concentration may be elevated because of an extracellular shift of potassium caused by insulin deficiency, hypertonicity, and acidemia. Patients with low-normal or low serum potassium concentration on admission have severe total-body potassium deficiency and require very

careful cardiac monitoring and more vigorous potassium replacement, because treatment lowers potassium further and can provoke cardiac dysrhythmia. The occurrence of stupor or coma in diabetic patients in the absence of definitive elevation of effective osmolality ( $\geq 320$  mOsm/kg) demands immediate consideration of other causes of mental status change. Effective osmolality may be calculated by the following formula:  $2[\text{measured Na (mEq/l)}] + \text{glucose (mg/dl)}/18$ . Amylase levels are elevated in the majority of patients with DKA, but this may be due to nonpancreatic sources, such as the parotid gland. A serum lipase determination may be beneficial in the differential diagnosis of pancreatitis. However, lipase could also be elevated in DKA. Abdominal pain and elevation of serum amylase and liver enzymes are noted more commonly in DKA than in HHS.

### Differential diagnosis

Not all patients with ketoacidosis have DKA. Starvation ketosis and alcoholic ketoacidosis (AKA) are distinguished by clinical history and by plasma glucose concentrations that range from mildly elevated (rarely  $>250$  mg/dl) to hypoglycemia. In addition, although AKA can result in profound acidosis, the serum bicarbonate concentration in starvation ketosis is usually not lower than 18 mEq/l. DKA must also be distinguished from other causes of high-anion gap metabolic acidosis, including lactic acidosis, ingestion of drugs such as salicylate, methanol, ethylene glycol, and paraldehyde, and chronic renal failure (which is more typically hyperchloremic acidosis rather than high-anion gap acidosis). Clinical history of previous drug intox-

ications or metformin use should be sought. Measurement of blood lactate, serum salicylate, and blood methanol level can be helpful in these situations. Ethylene glycol (antifreeze) is suggested by the presence of calcium oxalate and hippurate crystals in the urine. Paraldehyde ingestion is indicated by its characteristic strong odor on the breath. Because these intoxicants are low-molecular weight organic compounds, they can produce an osmolar gap in addition to the anion gap acidosis (14–16).

**TREATMENT** — Successful treatment of DKA and HHS requires correction of dehydration, hyperglycemia, and electrolyte imbalances; identification of comorbid precipitating events; and above all, frequent patient monitoring. Guidelines for the management of patients with DKA and HHS follow and are summarized in Figs. 1, 2, and 3. Table 3 includes a summary of major recommendations and evidence gradings.


### Fluid therapy

**Adult patients.** Initial fluid therapy is directed toward expansion of the intravascular and extravascular volume and restoration of renal perfusion. In the absence of cardiac compromise, isotonic saline (0.9% NaCl) is infused at a rate of  $15\text{--}20$  ml  $\cdot$  kg<sup>-1</sup> body wt  $\cdot$  h<sup>-1</sup> or greater during the 1st hour ( $\sim 1\text{--}1.5$  liters in the average adult). Subsequent choice for fluid replacement depends on the state of hydration, serum electrolyte levels, and urinary output. In general, 0.45% NaCl infused at  $4\text{--}14$  ml  $\cdot$  kg<sup>-1</sup>  $\cdot$  h<sup>-1</sup> is appropriate if the corrected serum sodium is normal or elevated; 0.9% NaCl at a similar rate is appropriate if corrected serum sodium is low. Once renal function is assured, the infusion should include 20–30 mEq/l potassium (2/3 KCl and 1/3 KPO<sub>4</sub>)

**Table 2—Typical total body deficits of water and electrolytes in DKA and HHS**

	DKA	HHS
Total water (liters)	6	9
Water (ml/kg)*	100	100–200
Na <sup>+</sup> (mEq/kg)	7–10	5–13
Cl <sup>-</sup> (mEq/kg)	3–5	5–15
K <sup>+</sup> (mEq/kg)	3–5	4–6
PO <sub>4</sub> (mmol/kg)	5–7	3–7
Mg <sup>2+</sup> (mEq/kg)	1–2	1–2
Ca <sup>2+</sup> (mEq/kg)	1–2	1–2

\*Per kilogram of body weight. †From Ennis et al. (15) and Kreisberg (8).



**Figure 1**—Protocol for the management of adult patients with DKA. \*DKA diagnostic criteria: blood glucose  $>250$  mg/dl, arterial pH  $<7.3$ , bicarbonate  $<15$  mEq/l, and moderate ketonuria or ketonemia. †After history and physical examination, obtain arterial blood gases; complete blood count with differential, urinalysis, blood glucose, blood urea nitrogen, electrolytes, chemistry profile, and creatinine levels STAT as well as an electrocardiogram. Obtain chest X ray and cultures as needed. ‡Serum Na should be corrected for hyperglycemia (for each 100 mg/dl glucose  $>100$  mg/dl, add 1.6 mEq to sodium value for corrected serum sodium value).

until the patient is stable and can tolerate oral supplementation. Successful progress with fluid replacement is judged by hemodynamic monitoring (improvement in blood pressure), measurement of fluid input/output, and clinical examination. Fluid replacement should correct estimated deficits within the first 24 h. The induced change in serum osmolality should not exceed  $3 \text{ mOsm} \cdot \text{kg}^{-1} \text{H}_2\text{O} \cdot \text{h}^{-1}$  (14–20, 22). In patients with renal or cardiac compromise, monitoring of serum osmolality and frequent assessment of cardiac, renal, and mental status must be performed during fluid resuscitation to avoid iatrogenic fluid overload (14–20,22).

**Pediatric patients (<20 years of age).** Initial fluid therapy is directed toward expansion of the intravascular and extravascular volume and restoration of renal perfusion. The need for vascular vol-


ume expansion must be offset by the risk of cerebral edema associated with rapid fluid administration. The 1st hour of fluids should be isotonic saline (0.9% NaCl) at the rate of  $10\text{--}20 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ . In a severely dehydrated patient, this may need to be repeated, but the initial reexpansion should not exceed  $50 \text{ ml/kg}$  over the first 4 h of therapy. Continued fluid therapy is calculated to replace the fluid deficit evenly over 48 h. In general, 0.9% NaCl infused at a rate of 1.5 times the 24-h maintenance requirements ( $\sim 5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ) will accomplish a smooth rehydration, with a decrease in osmolality not exceeding  $3 \text{ mOsm} \cdot \text{kg}^{-1} \text{H}_2\text{O} \cdot \text{h}^{-1}$ . Once renal function is assured and serum potassium is known, the infusion should include 20–40 mEq/l potassium (2/3 KCl and 1/3 KPO<sub>4</sub>). Once serum glucose reaches 250 mg/dl, fluid should be

changed to 5% dextrose and 0.45–0.75% NaCl, with potassium as described above. Therapy should include monitoring mental status to rapidly identify changes that might indicate iatrogenic fluid overload, which can lead to symptomatic cerebral edema (23–25).

### Insulin therapy

Unless the episode of DKA is mild (Table 1), regular insulin by continuous intravenous infusion is the treatment of choice. Once hypokalemia ( $\text{K}^+ < 3.3$  mEq/l) is excluded, an intravenous bolus of regular insulin at 0.15 U/kg body wt, followed by a continuous infusion of regular insulin at a dose of  $0.1 \text{ U} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$  (5–7 U/h in adults), should be administered. This low dose of insulin usually decreases plasma glucose concentration at a rate of  $50\text{--}75 \text{ mg} \cdot \text{dl}^{-1} \cdot \text{h}^{-1}$ , similar to a higher dose insulin

## Management of Adult Patients with HHS\*




**Figure 2**—Protocol for the management of adult patients with HHS. \*Diagnostic criteria: blood glucose >600 mg/dl, arterial pH >7.3, bicarbonate >15 mEq/L, effective serum osmolality >320 mOsm/kg H<sub>2</sub>O, and mild ketonuria or ketonemia. This protocol is for patients admitted with mental status change or severe dehydration who require admission to an intensive care unit. For less severe cases, see text for management guidelines. Effective serum osmolality calculation:  $2[\text{measured Na (mEq/L)}] + \text{glucose (mg/dl)}/18$ . †After history and physical examination, obtain arterial blood gases, complete blood count with differential, urinalysis, plasma glucose, blood urea nitrogen, electrolytes, chemistry profile, and creatinine levels STAT as well as an electrocardiogram. Chest X ray and cultures as needed. ‡Serum Na should be corrected for hyperglycemia (for each 100 mg/dl glucose >100 mg/dl, add 1.6 mEq to sodium value for corrected serum value).

regimen (26). If plasma glucose does not fall by 50 mg/dl from the initial value in the 1st hour, check hydration status; if acceptable, the insulin infusion may be doubled every hour until a steady glucose decline between 50 and 75 mg/h is achieved. When the plasma glucose reaches 250 mg/dl in DKA or 300 mg/dl in HHS, it may be possible to decrease the insulin infusion rate to  $0.05\text{--}0.1 \text{ U} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$  (3–6 U/h), and dextrose (5–10%) may be added to the intravenous fluids. Thereafter, the rate of insulin administration or the concentration of dextrose may need to be adjusted to maintain the above glucose values until acidosis in DKA or mental obtundation and hyperosmolality in HHS are resolved.

Ketonemia typically takes longer to clear than hyperglycemia. The nitroprusside method only measures acetoacetic acid and acetone. However,  $\beta$ -OHB, the strongest and most prevalent acid in DKA, is not measured by the nitroprusside method. During therapy,  $\beta$ -OHB is converted to acetoacetic acid, which may lead the clinician to believe that ketosis has worsened. Therefore, assessments of urinary or serum ketone levels by the nitroprusside method should not be used as an indicator of response to therapy. During therapy for DKA or HHS, blood should be drawn every 2–4 h for determination of serum electrolytes, glucose, blood urea nitrogen, creatinine, osmolality, and

venous pH (for DKA). Generally, repeat arterial blood gases are unnecessary; venous pH (which is usually 0.03 U lower than arterial pH) and anion gap can be followed to monitor resolution of acidosis. With mild DKA, regular insulin given either subcutaneously or intramuscularly every hour is as effective as intravenous administration in lowering blood glucose and ketone bodies (27). Patients with mild DKA should first receive a “priming” dose of regular insulin of 0.4–0.6 U/kg body wt, half as an intravenous bolus and half as a subcutaneous or intramuscular injection (22). Thereafter,  $0.1 \text{ U} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$  of regular insulin should be given subcutaneously or intramuscularly.

## Management of Pediatric Patients (&lt;20 years) with DKA\* or HHS†



**Figure 3**—Protocol for the management of pediatric patients (<20 years) with DKA or HHS. \*DKA diagnostic criteria: blood glucose >250 mg/dl, venous pH <7.3, bicarbonate <15 mEq/L, and moderate ketonuria or ketonemia. †HHS diagnostic criteria: blood glucose >600 mg/dl, venous pH >7.3, bicarbonate >15 mEq/L, and altered mental status or severe dehydration. ‡After the initial history and physical examination, obtain blood glucose, venous blood gases, electrolytes, blood urea nitrogen, creatinine, calcium, phosphorous, and urine analysis STAT. §Usually 1.5 times the 24-h maintenance requirements ( $\sim 5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ) will accomplish a smooth rehydration; do not exceed two times the maintenance requirement. ||The potassium in solution should be 1/3  $\text{KPO}_4$  and 2/3 KCl or Acetate.

After resolution of DKA (glucose <200 mg/dl, serum bicarbonate  $\geq 18$  mEq/L, venous pH >7.3, anion gap <12 mEq/L) and when patients are able to take fluids orally, a multidose regimen may be initiated based on history of previous treatment. However, for newly diagnosed patients, a total insulin dose of  $0.6\text{--}0.7 \text{ U} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$  may be initiated as a multidose regimen of short- and intermediate-/long-acting insulin, with subsequent modification based on glucose testing. Finally, some type 2 diabetic patients may be discharged on oral agents and dietary therapy.

### Potassium

Despite total-body potassium depletion, mild to moderate hyperkalemia is not uncommon in patients with hyperglycemic crises. Insulin therapy, correction of acidosis, and volume expansion decrease serum

potassium concentration. To prevent hypokalemia, potassium replacement is initiated after serum levels fall below 5.5 mEq/L, assuming the presence of adequate urine output. Generally, 20–30 mEq potassium (2/3 KCl and 1/3  $\text{KPO}_4$ ) in each liter of infusion fluid is sufficient to maintain a serum potassium concentration within the normal range of 4–5 mEq/L. Rarely, DKA patients may present with significant hypokalemia. In such cases, potassium replacement should begin with fluid therapy, and insulin treatment should be delayed until potassium concentration is restored to >3.3 mEq/L to avoid arrhythmias or cardiac arrest and respiratory muscle weakness.

### Bicarbonate

Bicarbonate use in DKA remains controversial (28). At a pH >7.0, reestablishing insulin activity blocks lipolysis and resolves

ketoacidosis without any added bicarbonate. Prospective randomized studies have failed to show either beneficial or deleterious changes in morbidity or mortality with bicarbonate therapy in DKA patients with pH between 6.9 and 7.1 (29). No prospective randomized studies concerning the use of bicarbonate in DKA with pH values <6.9 have been reported. Given that severe acidosis may lead to a myriad of adverse vascular effects, it seems prudent that for adult patients with a pH <6.9, 100 mmol sodium bicarbonate be added to 400 ml sterile water and given at a rate of 200 ml/h. In patients with a pH of 6.9–7.0, 50 mmol sodium bicarbonate is diluted in 200 ml sterile water and infused at a rate of 200 ml/h. No bicarbonate is necessary if pH is >7.0.

In the pediatric patient, there are no randomized studies in patients with pH <6.9. If the pH remains below 6.9 after the

Table 3—Summary of major recommendations

Recommendations	Grading
Initiate insulin therapy according to recommendations in position statement.	A
Unless the episode of DKA is mild, regular insulin by continuous intravenous infusion is preferred.	B
Assess need for bicarbonate therapy, and if necessary, follow treatment recommendations in position statement: Bicarbonate may be beneficial in patients with a pH <6.9; not necessary if pH is >7.0.	C
Studies have failed to show any beneficial effect of phosphate replacement on the clinical outcome in DKA. However, to avoid cardiac and skeletal muscle weakness and respiratory depression due to hypophosphatemia, careful phosphate replacement may sometimes be indicated in patients with cardiac dysfunction, anemia, or respiratory depression and in those with serum phosphate concentration <1.0 mg/dl.	A
Studies of cerebral edema in DKA are limited in number. Therefore, to avoid the occurrence of cerebral edema, follow the recommendations in the position statement regarding a gradual correction of glucose and osmolality as well as the judicious use of isotonic or hypotonic saline, depending on serum sodium and the hemodynamic status of the patient.	C
Initiate fluid replacement therapy based on recommendations in position statement.	A

Scientific evidence was ranked based on the American Diabetes Association's grading system. The highest ranking (A) is assigned when there is supportive evidence from well-conducted, generalizable, randomized controlled trials that are adequately powered, including evidence from a meta-analysis that incorporated quality ratings in the analysis. An intermediate ranking (B) is given to supportive evidence from well-conducted cohort studies, registries, or case-control studies. A lower rank (C) is assigned to evidence from uncontrolled or poorly controlled studies or when there is conflicting evidence with the weight of the evidence supporting the recommendation. Expert consensus (E) is indicated, as appropriate. For a more detailed description of this grading system, refer to *Diabetes Care* 24 (Suppl. 1):S83, 2001.

initial hour of hydration, it seems prudent to administer 1–2 mEq/kg sodium bicarbonate over an hour. This sodium bicarbonate can be added to 0.45 NaCl, with any required potassium, and this solution can be used as the rehydration solution for that hour. No bicarbonate therapy is required if pH is  $\geq 7.0$  (30,31).

Insulin, as well as bicarbonate therapy, lowers serum potassium; therefore, potassium supplementation should be maintained in intravenous fluid as described above and carefully monitored. (See Fig. 1 for guidelines.) Thereafter, venous pH should be assessed every 2 h until the pH rises to 7.0, and treatment should be repeated every 2 h if necessary. (See Kitabchi et al. [11] for a complete description of studies done to date on the use of bicarbonate in DKA.)

### Phosphate

Despite whole-body phosphate deficits in DKA that average  $\sim 1.0$  mmol/kg body wt, serum phosphate is often normal or increased at presentation. Phosphate concentration decreases with insulin therapy. Prospective randomized studies have failed to show any beneficial effect of phosphate

replacement on the clinical outcome in DKA (32), and overzealous phosphate therapy can cause severe hypocalcemia with no evidence of tetany (17,32). However, to avoid cardiac and skeletal muscle weakness and respiratory depression due to hypophosphatemia, careful phosphate replacement may sometimes be indicated in patients with cardiac dysfunction, anemia, or respiratory depression and in those with serum phosphate concentration <1.0 mg/dl. When needed, 20–30 mEq/l potassium phosphate can be added to replacement fluids. No studies are available on the use of phosphate in the treatment of HHS. Continuous monitoring using a flowsheet (Fig. 4) aids in the organization of recovery parameters and treatment interventions.

**COMPLICATIONS**— The most common complications of DKA and HHS include hypoglycemia due to overzealous treatment with insulin, hypokalemia due to insulin administration and treatment of acidosis with bicarbonate, and hyperglycemia secondary to interruption/discontinuation of intravenous insulin therapy after recovery without subsequent coverage with subcutaneous insulin. Com-

monly, patients recovering from DKA develop hyperchloremia caused by the use of excessive saline for fluid and electrolyte replacement and transient non-anion gap metabolic acidosis as chloride from intravenous fluids replaces ketoanions lost as sodium and potassium salts during osmotic diuresis. These biochemical abnormalities are transient and are not clinically significant except in cases of acute renal failure or extreme oliguria.

Cerebral edema is a rare but frequently fatal complication of DKA, occurring in 0.7–1.0% of children with DKA. It is most common in children with newly diagnosed diabetes, but has been reported in children with known diabetes and in young people in their twenties (33,34). Fatal cases of cerebral edema have also been reported with HHS. Clinically, cerebral edema is characterized by a deterioration in the level of consciousness, with lethargy, decrease in arousal, and headache. Neurological deterioration may be rapid, with seizures, incontinence, pupillary changes, bradycardia, and respiratory arrest. These symptoms progress as brain stem herniation occurs. The progression may be so rapid that papilledema is not found. Once the clinical symptoms other than lethargy and behavioral changes occur, mortality is high (>70%), with only 7–14% of patients recovering without permanent morbidity. Although the mechanism of cerebral edema is not known, it likely results from osmotically driven movement of water into the central nervous system when plasma osmolality declines too rapidly with the treatment of DKA or HHS. There is a lack of information on the morbidity associated with cerebral edema in adult patients; therefore, any recommendations for adult patients are clinical judgments, rather than scientific evidence. Prevention measures that might decrease the risk of cerebral edema in high-risk patients are gradual replacement of sodium and water deficits in patients who are hyperosmolar (maximal reduction in osmolality  $3 \text{ mOsm} \cdot \text{kg}^{-1} \text{ H}_2\text{O} \cdot \text{h}^{-1}$ ) and the addition of dextrose to the hydrating solution once blood glucose reaches 250 mg/dl. In HHS, a glucose level of 250–300 mg/dl should be maintained until hyperosmolarity and mental status improves and the patient becomes clinically stable (35).

Hypoxemia and, rarely, noncardiogenic pulmonary edema may complicate the treatment of DKA. Hypoxemia is attributed to a reduction in colloid osmotic pres-



- Clin Invest* 12:297–326, 1933
4. Halperin ML, Cheema-Dhadli S: Renal and hepatic aspects of ketoacidosis: a quantitative analysis based on energy turnover. *Diabetes Metab Rev* 5:321–336, 1989
  5. Malone ML, Gennis V, Goodwin JS: Characteristics of diabetic ketoacidosis in older versus younger adults. *J Am Geriatr Soc* 40:1100–1104, 1992
  6. Matz R: Hyperosmolar nonacidotic diabetes (HNAD). In *Diabetes Mellitus: Theory and Practice*. 5th ed. Porte DJ Jr, Sherwin RS, Eds. Amsterdam, Elsevier, 1997, p. 845–860
  7. Morris LE, Kitabchi AE: Coma in the diabetic. In *Diabetes Mellitus: Problems in Management*. Schnatz JD, Ed. Menlo Park, CA, Addison-Wesley, 1982, p. 234–251
  8. Kreisberg RA: Diabetic ketoacidosis: new concepts and trends in pathogenesis and treatment. *Ann Int Med* 88:681–695, 1978
  9. Klekamp J, Churchwell KB: Diabetic ketoacidosis in children: initial clinical assessment and treatment. *Pediatric Annals* 25:387–393, 1996
  10. Glaser NS, Kupperman N, Yee CK, Schwartz DL, Styne DM: Variation in the management of pediatric diabetic ketoacidosis by specialty training. *Arch Pediatr Adolescent Med* 151:1125–1132, 1997
  11. Kitabchi AE, Umpierrez GE, Murphy MB, Barrett EJ, Kreisberg RA, Malone JL, Wall BM: Management of hyperglycemic crises in patients with diabetes mellitus (Technical Review). *Diabetes Care* 24:131–153, 2001
  12. Beigelman PM: Severe diabetic ketoacidosis (diabetic coma): 482 episodes in 257 patients: experience of three years. *Diabetes* 20:490–500, 1971
  13. Polonsky WH, Anderson BJ, Lohrer PA, Aponte JE, Jacobson AM, Cole CF: Insulin omission in women with IDDM. *Diabetes Care* 17:1178–1185, 1994
  14. Kitabchi AE, Fisher JN, Murphy MB, Rumbak MJ: Diabetic ketoacidosis and the hyperglycemic hyperosmolar nonketotic state. In *Joslin's Diabetes Mellitus*. 13th ed. Kahn CR, Weir GC, Eds. Philadelphia, Lea & Febiger, 1994, p. 738–770
  15. Ennis ED, Stahl EJVB, Kreisberg RA: The hyperosmolar hyperglycemic syndrome. *Diabetes Rev* 2:115–126, 1994
  16. Marshall SM, Walker M, Alberti KGMM: Diabetic ketoacidosis and hyperglycaemic non-ketotic coma. In *International Textbook of Diabetes Mellitus*. 2nd ed. Alberti KGMM, Zimmet P, DeFronzo RA, Eds. New York, John Wiley, p. 1215–1229, 1997
  17. Carroll P, Matz R: Uncontrolled diabetes mellitus in adults: experience in treating diabetic ketoacidosis and hyperosmolar coma with low-dose insulin and uniform treatment regimen. *Diabetes Care* 6:579–585, 1983
  18. Ennis ED, Stahl EJ, Kreisberg RA: Diabetic ketoacidosis. In *Diabetes Mellitus: Theory and Practice*. 5th ed. Porte DJ Jr, Sherwin RS, Eds. Amsterdam, Elsevier, 1997, p. 827–844
  19. Hillman K: Fluid resuscitation in diabetic emergencies: a reappraisal. *Intensive Care Med* 13:4–8, 1987
  20. Fein IA, Rackow EC, Sprung CL, Grodman R: Relation of colloid osmotic pressure to arterial hypoxemia and cerebral edema during crystalloid volume loading of patients with diabetic ketoacidosis. *Ann Intern Med* 96:570–575, 1982
  21. Matz R: Hypothermia in diabetic acidosis. *Hormones* 3:36–41, 1972
  22. Kitabchi AE, Sacks HS, Young RT, Morris L: Diabetic ketoacidosis: reappraisal of therapeutic approach. *Ann Rev Med* 30:339–357, 1979
  23. Mahoney CP, Vleck BW, DelAguila M: Risk factors for developing brain herniation during diabetic ketoacidosis. *Pediatr Neurology* 21:721–727, 1999
  24. Finberg L: Why do patients with diabetic ketoacidosis have cerebral swelling, and why does treatment sometimes make it worse? *Pediatr Adolescent Med* 150:785–786, 1996
  25. Duck SC, Wyatt DT: Factors associated with brain herniation in the treatment of diabetic ketoacidosis. *J Pediatr* 113:10–14, 1988
  26. Kitabchi AE, Ayyagari V, Guerra SMO, Medical House Staff: The efficacy of low dose versus conventional therapy of insulin for treatment of diabetic ketoacidosis. *Ann Int Med* 84:633–638, 1976
  27. Fisher JN, Shahshahani MN, Kitabchi AE: Diabetic ketoacidosis: low dose insulin therapy by various routes. *N Engl J Med* 297:238–247, 1977
  28. Barnes HV, Cohen RD, Kitabchi AE, Murphy MB: When is bicarbonate appropriate in treating metabolic acidosis including diabetic ketoacidosis? In *Debates in Medicine*. Gitnick G, Barnes HV, Duffy TP, et al., Eds. Chicago, Yearbook, 1990, p. 172
  29. Morris LR, Murphy MB, Kitabchi AE: Bicarbonate therapy in severe diabetic ketoacidosis. *Ann Int Med* 105:836–840, 1986
  30. Vanelli M, Chiari G, Ghizzoni L, Costi G, Giacalone T, Chiarelli F: Effectiveness of a prevention program for diabetic ketoacidosis in children. *Diabetes Care* 22:7–9, 1999
  31. Viallon A, Zeni F, Lafond P, Venet C, Tardy B, Page Y, Bertrand JC: Does bicarbonate therapy improve the management of severe diabetic ketoacidosis? *Critical Care Medicine* 27, December 1999
  32. Fisher JN, Kitabchi AE: A randomized study of phosphate therapy in the treatment of diabetic ketoacidosis. *J Clin Endocrinol Metab* 57:177–180, 1983
  32. Rosenbloom AL: Intracerebral crises during treatment of diabetic ketoacidosis. *Diabetes Care* 13:22–33, 1990
  33. Duck SC, Wyatt DT: Factors associated with brain herniation in the treatment of diabetic ketoacidosis. *J Pediatr* 113:10–14, 1988
  34. Holsclaw DS Jr, Torcato B: Acute pulmonary edema in juvenile diabetic ketoacidosis. *Pediatr Pulmonology* 24:438–443, 1997
  35. Musey VC, Lee JK, Crawford R, Klatka MA, McAdams D, Phillips LS: Diabetes in urban African Americans. I. Cessation of insulin therapy is the major precipitating cause of diabetic ketoacidosis. *Diabetes Care* 18:483–489, 1995
  36. Umpierrez GE, Kelly JP, Navarrete JE, Casals MMC, Kitabchi AE: Hyperglycemic crises in urban blacks. *Arch Int Med* 157:669–675, 1997
  37. Fishbein HA, Palumbo PJ: Acute metabolic complications in diabetes. In *Diabetes in America*. National Diabetes Data Group, Ed. Bethesda, MD, National Institutes of Health, 1995, p. 283–291 (NIH pub. no. 95-1468)
  39. Kaufman FR, Halvorsen M: The treatment and prevention of diabetic ketoacidosis in children and adolescents with type 1 diabetes mellitus. *Pediatric Annals* 28:576–582, 1999